

## Galectin-1/LGALS1, Human, Recombinant

货号: PCK285

### 产品信息

别名 Beta-galactoside-binding lectin L-14-I, Galaptin, 14 kDa lectin, S-LAC lectin-1 Pricella

物种 Human

表达宿主 E.coli

序列信息 ACGLVASNLNLKPGECLRVRGEVAPDAKSFVLNLGKDSNNLCLHFNPRFN

> AHGDANTIVCNSKDGGAWGTEQREAVFPFQPGSVAEVCITFDQANLTVKL PDGYEFKFPNRLNLEAINYMAADGDFKIKCVAFD with polyhistidine tag at

the N-terminus.

检索号 P09382.2

分子量 15.5 kDa

标签 His-tag at the N-terminus

生物活性 Measured by its ability to agglutinate human red blood cells. The ED50 for this

effect is <2 μg/mL.

# 产品特性

>98% as determined by SDS-PAGE. Ni-NTA chromatography 纯度

内毒素 <0.1 EU per 1 µg of the protein by the LAL method.

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution,

> store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1%BSA,10%FBS,5%HSA or 5% trehalose solution),

protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH

7.4.

网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋








#### 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL.Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

The galectin-1 protein is 135 amino acids in length and highly conserved across species. It can be found in the nucleus, the cytoplasm, the cell surface and in the extracellular space. Galectins, in general, lack a traditional signal sequence but are still secreted across the plasma membrane. Although galectins in general, and Gal-1 in particular, were first described as lectins that bind βgalactosides, it is now clear from the literature that as well as being a lectin, Gal-1 is also engaged in many protein-protein interactions. Gal-1 plays a number of crucial roles in neuronal cell differentiation and survival in both the central and the peripheral nervous systems, and the establishment and maintenance of T-cell tolerance and homeostasis in vivo.

#### SDS-PAGE



网站: www.procell.com.cn 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋



