

## TGFβ2/ TGFB2, Human, Recombinant

货号：PCK092

### 产品信息

别名	Transforming Growth Factor beta-2; TGFB2; Polyergin; G-TSF; Glioblastoma-Derived T-cell suppressor factor; Cetermin; BSC-1 cell growth inhibitor; TGF-beta-2
物种	Human
表达宿主	Human Cells
序列信息	Ala303-Ser414
检索号	P61812
分子量	12.7 kDa
生物活性	Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 cells. The ED50 for this effect is 30-180pg/ml.

### 产品特性

纯度	>95% as determined by reducing SDS-PAGE.
内毒素	<1.0 EU per µg as determined by LAL test.
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 µm filtered solution of 4 mM HCl.




## 复融

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


## 背景介绍

Transforming Growth Factor beta-2 (TGF-β2) is a secreted protein which belongs to the TGF-beta family. It is known as a cytokine that performs many cellular functions and has a vital role during embryonic development. The precursor is cleaved into mature TGF-beta-2 and LAP, which remains non-covalently linked to mature TGF-beta-2 rendering it inactive. It is an extracellular glycosylated Protein. It is known to suppress the effects of Interleukin dependent T-cell tumors. Defects in TGFB2 may be a cause of non-syndromic aortic disease (NSAD).

## SDS-PAGE



## 生物活性



Measured by its ability to inhibit the IL-4-dependent proliferation of TF-1 cells. The ED50 for this effect is 30-180pg/ml.

