

## IL-1RA/ICIL-1RA/IRAP/IL-1RN, Human, Recombinant

货号 : PCK144

### 产品信息

别名	ICIL-1RA, IRAP, IL-1RN
物种	Human
表达宿主	E.coli
序列信息	MRPSGRKSSKMQAFRIWDVNQKTFYLRNNQLVAGYLQGPNVNLEEKIDVV PIEPHALFLGIHGGKMCLSCVKSGDETRLQLEAVNITDLSNRKQDKRFAFI RSDSGPTTSFESAACPGWFLCTAMEADQPVSLTNMPDEGVMVTKFYFQED E with polyhistidine tag at the C-terminus.
检索号	P18510
分子量	18.07 kDa
标签	His-tag at the C-terminus
生物活性	Measure by its ability to inhibit IL-1 alpha -dependent proliferation in D10.G4.1 cells. The ED50 for this effect is <50 ng/mL.

### 产品特性

纯度	>98% as determined by SDS-PAGE. Ni-NTA chromatography
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 µm filtered solution containing 1X PBS, pH 7.4.



## 复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100  $\mu\text{g/mL}$ . Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

## 背景介绍

The interleukin-1 receptor antagonist (IL-1RA) is a protein that in humans is encoded by the IL1RN gene. IL-1RA was initially called the IL-1 inhibitor and was discovered separately in 1984 by two independent laboratories. IL-1RA is an agent that binds non-productively to the cell surface interleukin-1 receptor (IL-1R), the same receptor that binds interleukin 1 (IL-1), preventing IL-1 from sending a signal to that cell.

## SDS-PAGE

