

IL-13/P600, Mouse, Recombinant

货号:PCK251

产品信息

别名	P600

- 物种 Mouse
- 表达宿主 E.coli
- 序列信息 MPVPRSVSLPLTLKELIEELSNITQDQTPLCNGSMVWSVDLAAGGFCVALD SLTNISNCNAIYRTQRILHGLCNRKAPTTVSSLPDTKIEVAHFITKLLSYTKQL FRHGPF with polyhistidine tag at the C-terminus.
- 检索号 P20109.1
- 分子量 13.06 kDa
- 标签 His-tag at the C-terminus
- 生物活性 Measure by its ability to induce TF-1 cells proliferation. The ED50 for this effect is <4 ng/mL. The specific activity of recombinant mouse IL-13 is > 2.5 x 10^5 IU/mg.

产品特性

纯度	>95% as determined by SDS-PAGE. Ni-NTA chromatography.	
内毒素	<0.1 EU per 1 µg of the protein by the LAL method.	
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g.0.1% BSA,10%FBS,5%HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.	
运输	Ambient temperature or ice pack.	
制剂	The protein was lyophilized from a $0.2 \mu m$ filtered solution containing 1X PBS, pH 7.4.	





复融

It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μ g/mL.Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

Interleukin 13 (IL-13) is a protein that in humans is encoded by the IL13 gene. IL-13 was first cloned in 1993 and is located on chromosome 5q31 with a length of 1.4kb. It has a mass of 13 kDa and folds into 4 alpha helical bundles. The secondary structural features of IL-13 are similar to that of Interleukin 4 (IL-4); however it only has 25% sequence homology to IL-4 and is capable of IL-4 independent signaling. IL-13 is a cytokine secreted by T helper type 2 (Th2) cells, CD4 cells, Natural killer T cell, Mast cell, Basophil cells, Eosinophil cells and Nuocyte cells. Interleukin-13 is a central regulator in IgE synthesis, goblet cell hyperplasia, mucus hypersecretion, airway hyperresponsiveness, fibrosis and chitinase up-regulation. It is a mediator of allergic inflammation and different diseases including asthma.

SDS-PAGE

kDa	М
75	
75	-
60	
45	-
35	-
25	-
15	-
10 www.	procell.com.c

